

ABSTRACT OF THE DISCLOSURE

Specific Cytokine Antagonists, including TNF antagonists and/or Interleukin-1 antagonists, are used as novel therapeutic agents for the treatment of hearing loss, including presbycusis and other forms of sensorineural hearing loss. The present invention provides a method for inhibiting the action of TNF and/or IL-1 antagonists for treating hearing loss in a human by administering a TNF antagonist and/or an IL-1 antagonist for reducing the inflammation affecting the auditory apparatus of said human, or for modulating the immune response affecting the auditory apparatus of said human, by administering a therapeutically effective dosage level to said human of a TNF antagonist and/or an IL-1 antagonist. Administration may be systemic, through the subcutaneous, intramuscular, oral, or intravenous routes; or by delivering an anatomically localized application in the region of the head. The TNF antagonist is selected from the group consisting of etanercept, infliximab, D2E7, CDP 571, or thalidomide; and the IL-1 antagonist is either IL-1 RA or IL-1R type II receptor. Antiviral agents may be added for treating certain patients.

REFERENCES

Adams et al. (1989), "Deafness, Dizziness, and Disorders of Equilibrium," Principles of Neurology Ch.14:226-246.

Arck, P.C., Troutt, A.B., Clark, D.A. (1997). Soluble receptors neutralizing TNF-a and IL-1 block stress-triggered murine abortion. Am J Reprod Immunol 37, 262-266.

Colotta F, Re F, Muzio M, Bertini R, Polentarutti N, Sironi M, Giri JG, Dower SK, Sims JE, Mantovani A. Interleukin-1 type II receptor: a decoy target for IL-1 that is regulated by IL-4. Science 1993;261:472-5.

Dinarello CA. Interleukin-1, interleukin-1 receptors and interleukin-1 receptor antagonist. International Reviews of Immunology 1998;16:457-99.

Eldman M, Brennan FM, Maini RN. The role of cytokines in rheumatoid arthritis. Ann Rev Immunol 1996;14:397.

Hefti (1994), "Neurotrophic Factor Therapy for Nervous System Degenerative Diseases," J. Neurobiol. 25:1418-1435.

Lovell DJ, Giannini EH, Reiff, A, et al. Etanercept in children with polyarticular juvenile rheumatoid arthritis. N Engl J Med 2000;342(11):763-9.

5 Murray, K.M., Dahl, S.L. (1997). Recombinant human tumor necrosis factor receptor (p75) fusion protein (TNFR:Fc) in rheumatoid arthritis. Ann Pharmacother 31, 1335-1338.

Mohler, K.M., Torrance, D.S., Smith, C.A., Goodwin, R.G., Stremmler, K.E., Fung, V.P., Madani, H., Widmer, M.B. (1993). Soluble tumor necrosis factor (TNF) receptors are effective therapeutic agents in lethal endotoxemia and function simultaneously as both TNF carriers and TNF antagonists. J Immunol 151, 1548-1561.

Nadol (1981) "The Aging Peripheral Hearing Mechanism," Aging: Communication Processes and Disorders Ch. 4:63-85.

Nadol (1993), "Hearing Loss," New England J. of Medicine 329:1092-1102.

Schuknecht (1974) Pathology of the Ear pp. 388-403.

20 Sims JE, Gayle MA, Slack JL, Alderson MR, Bird TA, Giri JG, Colotta F, Re F, Mantovani A, Shanebeck K, Grabstein KH, Dower SK. Interleukin 1 signaling occurs exclusively via the type I receptor. Proc Natl Acad Sci U S A 1993;90:6155-9.

Spoendlin (1984), "Primary Neurons and Synapses," Ultrastructural Atlas of the Inner Ear Ch. 6:133-164.

Wiley, S.R., Schooley, K., Smolak, P.J., Din, W.S., Huang, C.-P., Nicholl, J.K., Sutherland, G.R., Davis Smith, T., Rauch, C., Smith, C.A., Goodwin, R.G. (1995). Identification and characterization of a new member of the TNF family that induces apoptosis. Immunity 3, 673-682.

Wilson et al. (1980), "The Efficacy of Steroids in the Treatment of Idiopathic Sudden Hearing Loss," Arch Otolaryngol 106:772-776.

10 Wooley PH, Dutcher J, Widmer MB, et al. Influence of a recombinant human soluble tumor necrosis factor receptor FC fusion protein on type II collagen-induced arthritis in mice. J Immunol 1993;151:6602.